

AD_____

Award Number: W81XWH-11-1-0143

TITLE: SNF5 mutation leads to intractable pain in schwannomatosis patients

PRINCIPAL INVESTIGATOR: Steven G. Matsumoto

CONTRACTING ORGANIZATION: Oregon Health and Sciences University
Portland, OR 97239

REPORT DATE: July 2013

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE July 2013		2. REPORT TYPE Final		3. DATES COVERED 15 June 2011 – 14 June 2013	
4. TITLE AND SUBTITLE SNF5 mutation leads to intractable pain in schwannomatosis patients				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-11-1-0143	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Steven G. Matsumoto E-Mail: matsumos@ohsu.edu				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Oregon Health and Sciences University Portland, OR 97239				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Ablation of snf5, in vivo, results in an increase in the number and type of sensory neurons that express the capsaicin receptor, TRPV1. Using a culture system, we have determined that soluble factor is released by snf5-/- Schwann cells that acts on sensory neurons to induce the expression of TRPV1. This factor is greater than 10K molecular weight and does not affect neuron survival. The increase in the immunoreactivity of TRPV1 is correlated with an increase in the expression of functional capsaicin-sensitive ion fluxes.					
15. SUBJECT TERMS Capsaicin receptor expression, conditioned medium factor					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			USAMRMC
U	U	U	UU	11	19b. TELEPHONE NUMBER (include area code)

Table of Contents

	<u>Page</u>
Introduction.....	1
Body.....	2
Key Research Accomplishments.....	6
Reportable Outcomes.....	6
Conclusion.....	6
References.....	7

Introduction.

Schwannomatosis is a devastating disease that does not affect longevity but has profound effects on the patient's quality of life due to the presence of intractable pain. The cause of this pain is not known. Our hypothesis is that mutations in the SNF5 gene in Schwann cells and ganglionic satellite cells leads to enhanced pain sensitivity in peripheral sensory neurons (see Campana et al, 2007). Mutations in the human SNF5 gene are linked to schwannomatosis (Hulsebos et al., 2007; Boyd et al., 2008; Hadfield et al., 2008; Sestini et al., 2008; Patil et al., 2008). Mice homozygous for snf5 deletion are embryonic lethal while heterozygotes develop rhabdoid tumors and other malignancies (Roberts et al., 2000; Klochender et al., 2000). We are using a tamoxifen inducible Cre-mediated recombination system driven by a mouse proteolipid protein-1 (*Plp1*) promoter (Plp1-cre/ESR1; Jackson Labs) to target the KO to Schwann and satellite cells. When this mouse is crossed with a floxed snf5 mouse, gene activity is reduced by >80% in the peripheral nervous system. A description of our aims for this project and our preliminary data follows.

Project Proposed Aims and Results.

The first part of this report of results is essentially identical to our yearly progress report (**posted 2/12/2013**). To this belated annual report we add newly obtained data indicating a induction of capsaicin sensitivity by Schwann cell conditioned medium and the results of 2 microarrays.

We proposed 2 specific aims: 1. **Test the hypothesis that the loss of *snf5* increases pain sensitivity by increasing the expression of the capsaicin receptor (TRPV1) in polymodal nociceptors and by inducing capsaicin-sensitivity in sensory neurons of other modalities.**

We have utilized a tamoxifen-inducible *plpCre* / *snf5-flox/flox* mouse to produce an *in vivo* targeted deletion of SNF5. We are using the expression of the capsaicin receptor, TRPV1, to estimate the number of nociceptors in sensory ganglia of these animals.

In experiments where we acutely ablated *snf5* in adult mice, we detect an increase in the number of TRPV1-expressing neurons in the trigeminal and dorsal root ganglia (**figure 1**).




Figure 1. Trigeminal ganglion of oil injected (control; A,A') and tamoxifen injected (B, B') adult animals. Ganglia fixed 7 days after injections and labeled with anti-TRPV1. Panels are at 10X (A,B) and 20X (A', B').

To quantify the increase in TRPV1-IR neurons, we serially sectioned lumbar DRG (L1-5) and counted the neurons, \pm TRPV1-IR, in every third section. The average percent positive neurons in 5 ganglia/animal were compared using a student's t-test. The results of 4 animals (2 tamoxifen-injected and 2 oil injected controls) were significantly different, 68 ± 9 vs $42 \pm 6\%$ respectively; $p < 0.01$

Strikingly, in preliminary cell counts, we observe an increase in large diameter ($>24\mu\text{m}$; 47% in tamoxifen- vs 8% oil injected controls) TRPV1-IR sensory neurons in lumbar DRGs (not shown) and trigeminal ganglia (**figure 1**). Since the small diameter ($<20\mu\text{m}$) neurons are the typical nociceptors, this result suggests that sensory neurons of non-pain modalities are being recruited to express a nociceptor property. Consistent with this *in vivo* result, we detect an increase in large diameter, TRPV1-IR sensory neurons in cultures treated with conditioned medium (see figure 2 and Aim 2 below).

2. Test the hypothesis that the loss of *snf5* in Schwann and satellite cells results in an increase in their production and secretion of factors that increase the expression of TRPV1 in sensory neurons.

In our second model we used an *in vitro* sensory neuron system to screen the biological activity of soluble factors released into the medium by *snf5*-knockout Schwann cells. Dorsal root ganglion (DRG) neurons of neonate to postnatal day 18 animals were dissociated and plated at a density of approximately 1000 neurons / cm². The base culture medium consists of L15, modified for 5% CO₂ atmosphere, as described in Mains and Patterson, (1973) with 5% adult rat serum, antibiotics and 100 ng/ml of 7s NGF.

The base medium was supplemented with medium conditioned by Schwann cells (SC-CM) \pm *snf5* (ablated using cre-lentiviral infection). The SC-CM was collected from confluent Schwann cell cultures, and concentrated using Amicon Ultra centrifugal filters with a 10000 molecular weight cutoff (Millipore). The SC-CM \pm *snf5* was tested at 1-5 X concentration for 48-96 hours.




Figure 2. DRG culture treated with Control SC-CM (A) and -/- *snf5* SC-CM (B) for 48 hours and labeled with anti-TRPV1. Note the presence of large diameter TRPV1-IR neurons in (B)

In these experiments, dissociated neurons were plated onto laminin-coated, 1 cm diameter glass coverslips. After CM treatment, the cultures were fixed and labeled with anti-TRPV1 antibodies and the appropriate secondary antibodies. For cell counts, each coverslip was divided into 4 sections and 3 randomly selected 20X fields were counted.

After 48 hours in CM we observe an increase in the percentage of TRPV1-IR neurons in cultures treated with -*snf5* SC-CM ($73 \pm 7\%$) vs control SC-CM treatment ($52 \pm 2\%$). In addition, we observe both large and small diameter TRPV1-IR neurons in the *snf5* -/-, SC-CM treatment but predominately small diameter neurons in the control SC-control CM, as we observed in the sensory ganglia *in situ* (not quantified at this time, figure 1)

We used a histochemical assay to determine the capsaicin sensitivity of sensory neurons cultured with \pm *SNF5* SC-CM. In this assay, DRG cultures were incubated in a saline solution containing 0 calcium and 5 mM cobalt \pm capsaicin (50 μ M). After washing, intracellular cobalt was precipitated with ammonium sulfide. Following fixation, the cobalt precipitate was enhanced with silver using the Timm's intensification protocol (Matsumoto, 1994). Using this method, we detect a significant increase ($p < .01$) in capsaicin-sensitive neurons in -*snf5* SC-CM ($94 \pm 1\%$) vs $58 \pm 2\%$ in control SC-CM treated cultures (figure 3)




Figure 3. DRG culture, control SC-CM (A) and -snf5 SC-CM (B) treated with 50 μ M capsaicin in the presence of 5 mM cobalt and subsequently enhanced with Timm's silver intensification protocol. Note, small diameter neurons labeled in (*) in A) while many large and small diameter neurons labeled in (B).

Capsaicin Sensitivity can be Rapidly Enhanced by SC-CM. In a second series of culture experiments, we determined that the sensitivity of dorsal root ganglion neurons (as measured by cobalt uptake) increased after a 1 hour exposure to SNF5 $-/-$ conditioned CM. Control cultures displayed 56% positive neurons vs 82% in CM-treated cultures ($p < 0.001$). This increase in capsaicin-sensitive neurons occurred in the absence of increased TRPV1-immunoreactivity.




Figure 4. DRG culture treated with control SC-CM (left panel) and snf5SC-CM (right panel) for 1 hour. Both cultures were exposed to 50 μ M capsaicin in the presence of 5 mM cobalt and subsequently enhanced with Timm's silver intensification protocol as previously described.

Microarray analysis of Schwann cells \pm snf5 and DRG sensory neurons treated with control and snf5 $-/-$ Schwann cell conditioned medium. Schwann cells from snf5 double-floxed animals were prepared in the following manner:

Sciatic nerves and brachial plexuses from P2-P5 snf5 $-/-$ double-floxed pups were dissected and dissociated with collagenase and trypsin treatment. Schwann cells are cultured in DMEM supplemented with 10% FBS, 100U/ml penicillin/streptomycin, 2ng/ml human heregulin- β 1, 0.5 μ M forskolin and 10ng/ml human bFGF.

To ablate snf5, the cultured cells were infected with either control or Cre- lentivirus. For the infection; 2×10^5 Schwann cells were plated on D-polylysine coated 10 cm culture dish one day before infection. About 20×10^5 transforming units (TU) were used for the infection (MOI 1:5-1:10). Culture medium was changed the next morning and the cells were maintained for another 2 to 3 days before collecting RNA.

The analysis of the microarrays data shows that the SNF5 does not affect the cell cycle. This result is in concordance with our in vitro data. The analysis does show an up regulation of several matrix metallopeptidases that are known to affect the development of tumors and tumor progression. Among these MMP 12, MMP 1a and MMP 10.

MMP 12 up 2.05

MMP 1a up 2.02

MMP 10 up 1.96

MMP 9 up 1.56

Our data also show an up-regulation of certain Chemokines /Chemokines receptors such as CCL5 (up 1.54) and CCR1(up 1.51).

We are continuing to analyze this data and identify possible links to our pain phenotype of enhanced capsaicin sensitivity,

Microarray of DRG sensory neurons. Wildtype DRG's were harvested from P7 pups and dissociated with collagenase/dispase. The neurons were preplated on uncoated culture plastic for 2 hours to reduce non-neuronal cells. The cells were gently resuspended and plated on to polylysine/laminin treated dishes. The culture medium contained 10 μ M fUdr and uracil for 3 days. The neuron enriched cultures were treated with control or snf5-/- SC-CM for 24 hours and RNA collected.

The result of the neuron microarray is as follows:

Select transcripts up-regulated in DRG sensory neurons treated with CM from SNF5 -/-:

Transcripts	Fold change
RTP4 a Golgi chaperone (protects mu-delta receptors from ubiquitination and degradation).	1.7
ISG15 a tumor suppressor	1.84
Z-DNA binding protein1 role in the transport and translation of mRNA in axonal regeneration	1.55
Calcium channel voltage dependant alpha-2/delta subunit (Cacna2d1): Up regulated in neuropathic pain. Targeted by drugs used for focal epilepsy, neuropathic pain and anxiety disorders	1.35
Mu opioid receptors variant A	1.3
Sodium channel 207=Nav1.2	1.2
Glutamate receptor ionotropic AMPA2 (alpha2)	1.23

Among transcripts down-regulated in DRG sensory neurons treated with CM from SNF5 -/-:

4

Transcripts	Fold change
Ras homolog gene family, member A (Rhoa)	1.61
Poly(A) binding protein, cytoplasmic 4-like (Pabpc4l),	1.54
Versican (Vcan) Variant 2 (inhibit cell proliferation and tumor growth)	1.7
DNA cytosine methyl transferase	1.99

As in the case of the Schwann cell array, we are continuing to analyze this data and identify possible links to our pain phenotype of enhanced capsaicin sensitivity,

Key Research Accomplishments.

Target-deletion of SNF5 leads to increase TRPV1 expression in sensory neurons

SNF5 ^{-/-} Schwann cells secrete a factor that stimulates the expression of TRPV1

The conditioned medium factor increases the capsaicin sensitivity of multiple classes of sensory neuron.

Conditioned medium from snf5^{-/-} Schwann cells has long term effects (i.e. increased TRPV1 expression) and short term effects (i.e. rapid increase in capsaicin sensitivity). In the short term experiments, we did not detect a significant increase in the number of TRPV1-IR neurons in cultures that demonstrated increased capsaicin-sensitivity. This may be due to a lack of sensitivity in our immunocytochemistry or our inability to detect a redistribution of TRPV1 receptors (e.g. Zhang et al, 2005)

Reportable Outcomes.

Manuscript in preparation

Conclusion.

The preliminary identification of a conditioned medium factor (CM) that is >10K mw is interesting and potentially very important towards advancing our understanding of this disease. Our finding that the CM factor induces TRPV1 expression in large diameter sensory neurons is significant, since it suggests that one outcome of the mutation may be to convert non-pain sensory neurons to this phenotype.

Our microarray analysis of Schwann cells \pm snf5 and DRG sensory neurons treated with Schwann cell condition medium has identified potential secreted Schwann cell-derived factors and neuronal target genes.

References.

Boyd C, Smith M, Kluwe L, Balogh A, Maccollin M, Plotkin S. Alterations in the SMARCB1 (INI1) tumor suppressor gene in familial schwannomatosis. *Clin Genet*. 2008 Jul 21.

Campana WM. Schwann cells: activated peripheral glia and their role in neuropathic pain. *Brain Behav Immun*. 2007 Jul;21(5):522-7.

Hulsebos TJ, Plomp AS, Wolterman RA, Robanus-Maandag EC, Baas F, Wesseling P. Germline mutation of INI1/SMARCB1 in familial schwannomatosis. *Am J Hum Genet*. 2007 Apr;80(4):805-10.

Hadfield, KD, Newman, WG, Bowers, NL, Wallace, A, Bolger, C, Colley A, McCann, E, Trump D, Prescott T Evans, DG Molecular characterization of SMARCB1 and NF2 in familial and sporadic schwannomatosis. *J. Med. Genet*. 2008Jun;45(9):608

Klochender-Yeivin A, Fiette L, Barra J, Muchardt C, Babinet C, Yaniv M. The murine SNF5/INI1 chromatin remodeling factor is essential for embryonic development and tumor suppression.

Page: 13

Mains RE, Patterson, PH Primary cultures of dissociated sympathetic neurons. I Establishment of long-term growth in culture and studies of differentiated properties. 1973 *J. Cell Biol*. Nov; 59:329

Matsumoto, SG Neuronal differentiation in cultures of murine neural crest II. Development of capsaicin-sensitive neurons. 1994 *Brain Res. Dev. Brain Res*. Nov 83: 17-27

Patil S, Perry A, Maccollin M, Dong S, Betensky RA, Yeh TH, Gutmann DH, Stemmer-Rachamimov AO. Immunohistochemical Analysis Supports a Role for INI1/SMARCB1 in Hereditary Forms of Schwannomas, but Not in Solitary, Sporadic Schwannomas. *Brain Pathol*. 2008 Apr 15.

Roberts CW, Galusha SA, McMenamin ME, Fletcher CD, Orkin SH. Haploinsufficiency of Snf5 (integrator interactor 1) predisposes to malignant rhabdoid tumors in mice. *Proc Natl Acad Sci U S A*. 2000 Dec 5;97(25):13796-800.

Sestini R, Bacci C, Provenzano A, Genuardi M, Papi L. Evidence of a four-hit mechanism involving SMARCB1 and NF2 in schwannomatosis-associated schwannomas. *Hum Mutat*. 2008 Feb;29(2):227-31.

Zhang X, Huang J, McNaughton PA NGF rapidly increases membrane expression of TRPV1 heat-gated ion channels. *EMBO* 2005 24: 4211-4223

